Emissions of Greenhouse Gases in the United States 2004

Report #: DOE/EIA-0573(2004)
Released Date: December 2005
Next Release Date: December 2006

Table 4. Numerical Estimates of Global Warming Potentials Compared With Carbon Dioxide
(Kilogram of Gas per Kilogram of Carbon Dioxide)

Gas	Lifetime (Years)	Direct Effect for Time Horizons of		
		20 Years	100 Years	500 Years
Carbon Dioxide	5 – 200[a]	1	1	1
Methane	12	62	23	7
Nitrous Oxide	114	275	296	156
HFCs, PFCs, and Sulfur Hexafluoride				
HFC-23	260	9,400	12,000	10,000
HFC-125	29	5,900	3,400	1,100
HFC-134a	13.8	3,300	1,300	400
HFC-152a	1.4	410	120	37
HFC-227ea	33	5,600	3,500	1,100
Perfluoromethane (CF_4)	50,000	3,900	5,700	8,900
Perfluoroethane (C_2F_6)	10,000	8,000	11,900	18,000
Sulfur Hexafluoride (SF_6)	3,200	15,100	22,200	32,400

[a]No single lifetime can be defined for carbon dioxide due to different rates of uptake by different removal processes.

Note: The typical uncertainty for global warming potentials is estimated by the Intergovernmental Panel on Climate Change at ±35 percent.

Source: Intergovernmental Panel on Climate Change, *Climate Change 2001: The Scientific Basis* (Cambridge, UK: Cambridge University Press, 2001), pp. 38 and 388-389.