

Table 26. Gas Home Customer-Weighted Heating Degree Days

Census Divisions	Cumulative November 1 through November 30				
	Normal ^a	1998	1999	Percent Change	
				Normal to 1999	1998 to 1999
New England CT, ME, MA, NH, RI, VT	693	711	608	-12.3	-14.5
Middle Atlantic NJ, NY, PA	646	618	536	-17.0	-13.3
East North Central IL, IN, MI, OH, WI	730	642	592	-18.9	-7.8
West North Central IA, KS, MN, MO, ND, NE, SD	788	673	564	-28.4	-16.2
South Atlantic DE, FL, GA, MD and DC, NC, SC, VA, WV	421	391	358	-15.0	-8.4
East South Central AL, KY, MS, TN	431	362	350	-18.8	-3.3
West South Central AR, LA, OK, TX	280	214	197	-29.6	-7.9
Mountain AZ, CO, ID, MT, NV, NM, UT, WY	715	651	546	-23.6	-16.1
Pacific ^b CA, OR, WA	341	382	309	-9.4	-19.1
U.S. Average ^b	559	514	452	-19.1	-12.1

^a Normal is based on calculations of data from 1961 through 1990.

^b Excludes Alaska and Hawaii.

Note: See Appendix A, Explanatory Note

10 for discussion of Heating Degree-Days computations.

Sources: National Oceanic and Atmospheric Administration.